

AD-A120 395 FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1
ASSOCIATION OF NORMAL RANDOM VARIABLES AND SLEPIAN'S INEQUALITY--ETC (U)
APR 82 K JOGDEO, M D PEARLMAN, L D PITT F49620-82-K-0007
UNCLASSIFIED FSU-STATISTICS-M620 AFOSR-TR-82-0899 NL

AD-A120 395 FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1-
ASSOCIATION OF NORMAL RANDOM VARIABLES AND SLEPIAN'S INEQUALITY--ETC (U)
APR 82 K JOGDEO, M D PEARLMAN, L D PITT F49620-82-K-0007
UNCLASSIFIED FSU-STATISTICS-M620 AFOSR-TR-82-0899 NL

AD-A120 395 FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1-
ASSOCIATION OF NORMAL RANDOM VARIABLES AND SLEPIAN'S INEQUALITY--ETC (U)
APR 82 K JOGDEO, M D PEARLMAN, L D PITT F49620-82-K-0007
UNCLASSIFIED FSU-STATISTICS-M620 AFOSR-TR-82-0899 NL

AD-A120 395 FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1-
ASSOCIATION OF NORMAL RANDOM VARIABLES AND SLEPIAN'S INEQUALITY--ETC (U)
APR 82 K JOGDEO, M D PEARLMAN, L D PITT F49620-82-K-0007
UNCLASSIFIED FSU-STATISTICS-M620 AFOSR-TR-82-0899 NL

AD-A120 395 FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1-
ASSOCIATION OF NORMAL RANDOM VARIABLES AND SLEPIAN'S INEQUALITY--ETC (U)
APR 82 K JOGDEO, M D PEARLMAN, L D PITT F49620-82-K-0007
UNCLASSIFIED FSU-STATISTICS-M620 AFOSR-TR-82-0899 NL

AD-A120 395 FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1-
ASSOCIATION OF NORMAL RANDOM VARIABLES AND SLEPIAN'S INEQUALITY--ETC (U)
APR 82 K JOGDEO, M D PEARLMAN, L D PITT F49620-82-K-0007
UNCLASSIFIED FSU-STATISTICS-M620 AFOSR-TR-82-0899 NL

AD-A120 395 FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1-
ASSOCIATION OF NORMAL RANDOM VARIABLES AND SLEPIAN'S INEQUALITY--ETC (U)
APR 82 K JOGDEO, M D PEARLMAN, L D PITT F49620-82-K-0007
UNCLASSIFIED FSU-STATISTICS-M620 AFOSR-TR-82-0899 NL

AD-A120 395 FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1-
ASSOCIATION OF NORMAL RANDOM VARIABLES AND SLEPIAN'S INEQUALITY--ETC (U)
APR 82 K JOGDEO, M D PEARLMAN, L D PITT F49620-82-K-0007
UNCLASSIFIED FSU-STATISTICS-M620 AFOSR-TR-82-0899 NL

AD-A120 395 FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1-
ASSOCIATION OF NORMAL RANDOM VARIABLES AND SLEPIAN'S INEQUALITY--ETC (U)
APR 82 K JOGDEO, M D PEARLMAN, L D PITT F49620-82-K-0007
UNCLASSIFIED FSU-STATISTICS-M620 AFOSR-TR-82-0899 NL

④

101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532
 533
 534
 535
 536
 537
 538
 539
 540
 541
 542
 543
 544
 545
 546
 547
 548
 549
 550
 551
 552
 553
 554
 555
 556
 557
 558
 559
 560
 561
 562
 563
 564
 565
 566
 567
 568
 569
 570
 571
 572
 573
 574
 575
 576
 577
 578
 579
 580
 581
 582
 583
 584
 585
 586
 587
 588
 589
 590
 591
 592
 593
 594
 595
 596
 597
 598
 599
 600
 601
 602
 603
 604
 605
 606
 607
 608
 609
 610
 611
 612

DATE
FILMED

1182

D1 6

AD A120395

Association of Normal Random Variables
and Slepian's Inequality

Abbreviated Title
Association of Normal Variables

by

Kumar Jogdeo
University of Illinois at Urbana-Champaign
Department of Mathematics
Urbana, Illinois 61801

Michael D. Perlman
University of Washington
Department of Statistics
Seattle, Washington 98195

Loren D. Pitt
University of Virginia
Department of Mathematics
Charlottesville, Virginia 22903

April, 1982
FSU Statistics Report No. M620
AFOSR Report No. 82-144

Department of Statistics
The Florida State University
Tallahassee, Florida 32306

DTIC
ELECTE
OCT 18 1982
S D
B

AMS 1970 subject classifications: Primary 62H05, Secondary 62N05.

Key Words and Phrases: normal variables, association, Slepian's inequality.

Research supported by Air Force Office of Scientific Research,
U.S.A.F., under Contract No. F49620-82-K-0007.

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

82 10 18 046

DTIC FILE COPY

I. Introduction

We give a simple proof of the result in Pitt (1981) that positively correlated normal random variables are associated. The proof is an adaptation of the original proof of Slepian's inequality in Slepian (1962), and extends to the case of elliptically contoured distributions.

II. Normal Random Vectors

Let $X = (x_1, \dots, x_n)$ be a mean zero n -dimensional normal random vector with $n \times n$ covariance matrix $\Sigma = (\sigma_{ij})$. By a smooth function we will mean a C^2 function $h(x)$ which together with its first and second order derivatives satisfy a $O(|x|^N)$ growth condition at ∞ , for some finite N .

We set

$$H(\Sigma) = Eh(X),$$

and we are interested in the manner that $H(\Sigma)$ varies with Σ . Our main result is

Proposition 1: Let Γ be another covariance matrix with $\gamma_{ii} = \sigma_{ii}$ and $\gamma_{ij} \leq \sigma_{ij}$ for all i and j . If h is a smooth function on R^n and if

$$(1) \quad \frac{\partial^2 h(x)}{\partial x_i \partial x_j} \geq 0 \quad \text{for all } i \text{ and } j \text{ with } \gamma_{ij} < \sigma_{ij},$$

then

$$(2) \quad H(\Gamma) \leq H(\Sigma).$$

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)

NOTICE OF TRANSMITTAL TO DTIC

This technical report has been reviewed and is approved for public release IAW AFR 190-12. Distribution is unlimited.

MATTHEW J. KERPER

Chief, Technical Information Division



Availability Codes	
Dist	Avail and/or Special
A	

Proof: By standard approximation arguments it suffices to show

$$\frac{\partial H(\Sigma)}{\partial \sigma_{ij}} \geq 0$$

whenever Σ is nonsingular and $\partial^2 h / \partial x_i \partial x_j \geq 0$. Let $\phi(x) = \phi_{\Sigma}(x)$ be the mean zero normal density on R^n with covariance matrix Σ . Then

$$(3) \quad \frac{\partial \phi}{\partial \sigma_{ii}} = \frac{1}{2} \frac{\partial^2 \phi}{\partial x_i^2}, \quad \frac{\partial \phi}{\partial \sigma_{ij}} = \frac{\partial^2 \phi}{\partial x_i \partial x_j}, \quad i \neq j.$$

See e.g. Plackett (1954). Using (3) and our assumptions on h which justify two integrations by parts we have

$$\begin{aligned} (4) \quad \frac{\partial H(\Sigma)}{\partial \sigma_{ij}} &= \int_{R^n} h(x) \frac{\partial \phi(x)}{\partial \sigma_{ij}} dx \\ &= \int_{R^n} \frac{\partial^2 h(x)}{\partial x_i \partial x_j} \phi(x) dx \\ &\geq 0, \end{aligned}$$

which completes the proof.

By varying h , Γ and Σ we obtain other results.

Corollary 1. Let $h(x_1, \dots, x_n) = f(x_1, \dots, x_k)g(x_{k+1}, \dots, x_n)$ where f and g are bounded measurable increasing functions. Suppose also that

$$\begin{aligned} \gamma_{ii} &= \sigma_{ii} && \text{for all } i, \\ \gamma_{ij} &= \sigma_{ij} && \text{if } 1 \leq i, j \leq k \text{ or } k < i, j \leq n, \\ \gamma_{ij} &\leq \sigma_{ij} && \text{if } 1 \leq i \leq k < j \leq n. \end{aligned}$$

Then

$$H(\Gamma) \leq H(\Sigma).$$

In particular, if $\sigma_{ij} \geq 0$ for $1 \leq i \leq k < j \leq n$.

$$Ef(x_1, \dots, x_k)Eg(x_{k+1}, \dots, x_n) \leq Ef(x_1, \dots, x_k)g(x_{k+1}, \dots, x_n).$$

Proof: If f and g are smooth then h is smooth and

$$\frac{\partial^2 h}{\partial x_i \partial x_j} = \frac{\partial f}{\partial x_i} \frac{\partial g}{\partial x_j} \geq 0 \quad \text{for } 1 \leq i \leq k < j \leq n.$$

In this case the result follows from (2). The general case follows by approximations as in Pitt (1981).

To state Slepian's inequality, we write $P_\Sigma(A)$ for the probability of the event that $X \in A \subset \mathbb{R}^n$.

Corollary 2. (Slepian) If $\gamma_{ii} = \sigma_{ii}$ and $\gamma_{ij} \leq \sigma_{ij}$ for all i and j then for each λ

$$P_\Gamma\{\max X_i \leq \lambda\} \leq P_\Sigma\{\max X_i \leq \lambda\}.$$

Proof: Again by standard arguments it suffices to show that $H(\Gamma) \leq H(\Sigma)$ for each product $h(x) = \prod_1^n f_i(x_i)$ of bounded non-negative smooth decreasing function $f_i(x_i)$. For $i \neq j$, each such product satisfies $\partial^2 h / \partial x_i \partial x_j \geq 0$, and the result follows from Proposition 1.

Remark: Several variants are possible. In particular, changing the sign of x_1, \dots, x_k in Corollary 1 gives the result of Jogdeo and Proschan (1981):

If $\sigma_{ij} \leq 0$ for $1 \leq i \leq k < j < n$, and if f and g are increasing then

$$Ef(x_1, \dots, x_k)g(x_{k+1}, \dots, x_n) \leq Ef(x_1, \dots, x_k)Eg(x_{k+1}, \dots, x_n).$$

III. Elliptically Contoured Distributions

The previous results extend to elliptically contoured distributions. The extension of Slepian's inequality to this case was given in Das Gupta, Eaton, Olkin, Perlman, Savage and Sobel (1972).

Let $\langle x, y \rangle$ denote the Euclidean inner product on R^n and let Σ be a non-singular positive definite matrix. A probability density on R^n of the form

$$p_{\Sigma}(x) = |\Sigma|^{-1/2} p(\langle x, \Sigma^{-1}x \rangle)$$

is called elliptically contoured. Here $p(\lambda) \geq 0$ is defined on $[0, \infty)$ and is assumed to satisfy $\int_0^{\infty} \lambda^{n-1} p(\lambda) d\lambda < \infty$.

We write

$$H(\Sigma) = \int_{R^n} h(x) p_{\Sigma}(x) dx,$$

and with this notation we will show that Proposition 1 remains valid.

It will suffice to establish (2) under the technical condition that $p(\lambda)$ is a C^2 function with compact support, and under this hypothesis we will show that:

$$(5) \quad \text{If } \partial^2 h / \partial x_i \partial x_j \geq 0, \text{ then } \frac{\partial}{\partial \sigma_{ij}} H(\Sigma) \geq 0.$$

The proof is similar to the earlier one, but requires a substitute for the equations (3). This is supplied by Proposition 2.

For $\lambda \geq 0$ we set

$$F(\lambda) = \int_0^\lambda p(\xi) d\xi.$$

Let

$$F_\infty = \int_0^\infty p(\xi) d\xi,$$

and

$$G_\Sigma(x) = (2|\Sigma|)^{-1/2} (F_\infty - F(\langle x, \Sigma^{-1} x \rangle)).$$

Proposition 2. If $h(x)$ is smooth and $p(\lambda)$ is C^2 with compact support, then

$$(6) \quad \frac{\partial}{\partial \sigma_{ij}} \int_{\mathbb{R}^n} h(x) p_\Sigma(x) dx = \int_{\mathbb{R}^n} \frac{\partial^2 h(x)}{\partial x_i \partial x_j} G_\Sigma(x) dx.$$

Remarks: Since $G_\Sigma \geq 0$ this proves (5). Also, in the case that $p_\Sigma = \phi_\Sigma$ is a normal density one easily checks that $G_\Sigma = \phi_\Sigma$, so (6) is a generalization of (3).

Proof: Let $\Sigma^{-1} = (\sigma^{ij})$. We will use the matrix identities

$$(7) \quad \frac{\partial}{\partial \sigma_{ii}} |\Sigma|^{-1/2} = -\frac{1}{2} \sigma^{ii} |\Sigma|^{-1/2},$$

$$\frac{\partial}{\partial \sigma_{ij}} |\Sigma|^{-1/2} = -\sigma^{ij} |\Sigma|^{-1/2}, \quad i \neq j$$

$$\frac{\partial}{\partial \sigma_{ii}} \langle x, \Sigma^{-1} x \rangle = -\left(\sum_{k=1}^n \sigma^{ik} x_k \right)^2,$$

$$\frac{\partial}{\partial \sigma_{ij}} \langle x, \Sigma^{-1} x \rangle = -2 \left(\sum_{k=1}^n \sigma^{ik} x_k \right) \left(\sum_{\ell=1}^n \sigma^{j\ell} x_\ell \right), \quad i \neq j.$$

without further comment.

Calculating, we now have

$$\begin{aligned}\frac{\partial p_{\Sigma}}{\partial \sigma_{ij}} &= -\sigma^{ij} p_{\Sigma} - 2|\Sigma|^{-\frac{1}{2}} p'(\langle x, \Sigma^{-1} x \rangle) \left(\sum_{k=1}^n \sigma^{ik} x_k \right) \left(\sum_{\ell=1}^n \sigma^{j\ell} x_{\ell} \right) \\ &= -\sigma^{ij} p_{\Sigma} - \left(\sum_{k=1}^n \sigma^{ik} x_k \right) \frac{\partial p_{\Sigma}}{\partial x_j}.\end{aligned}$$

Our assumptions on p justify integration by parts and we have

$$\begin{aligned}\frac{\partial}{\partial \sigma_{ij}} \int_{\mathbb{R}^n} h(x) p_{\Sigma}(x) dx &= -\sigma^{ij} \int_{\mathbb{R}^n} h(x) p_{\Sigma}(x) dx + \int_{\mathbb{R}^n} \frac{\partial}{\partial x_j} \left[\left(\sum_{k=1}^n \sigma^{ik} x_k \right) h(x) \right] p_{\Sigma}(x) dx \\ &= \int_{\mathbb{R}^n} \left(\sum_{k=1}^n \sigma^{ik} x_k \right) \frac{\partial h(x)}{\partial x_j} p_{\Sigma}(x) dx \\ &= \frac{1}{2|\Sigma|^{\frac{1}{2}}} \int_{\mathbb{R}^n} \frac{\partial h(x)}{\partial x_j} \frac{\partial}{\partial x_i} F(\langle x, \Sigma^{-1} x \rangle) dx \\ &= \int_{\mathbb{R}^n} \frac{\partial^2 h(x)}{\partial x_i \partial x_j} G_{\Sigma}(x) dx.\end{aligned}$$

The above calculations allow one extension which is perhaps worthwhile making. Using (7) we can easily verify that

$$(8) \quad \frac{\partial}{\partial \sigma_{ii}} \int_{\mathbb{R}^n} h(x) p_{\Sigma}(x) dx = \frac{1}{2} \int_{\mathbb{R}^n} \frac{\partial^2 h(x)}{\partial x_i^2} G_{\Sigma}(x) dx.$$

Combining (6) and (8) it is elementary to prove

Proposition 3. Let Γ and Σ be positive definite matrices and set

$A = \Sigma - \Gamma = (a_{ij})$. Let $\Sigma_t = \Gamma + tA$ and let $h(x)$ be a smooth function satisfying

$$(9) \quad Ah(x) \equiv \sum_{i,j=1}^n a_{ij} \frac{\partial^2 h(x)}{\partial x_i \partial x_j} \geq 0.$$

Then

$$\int_{\mathbb{R}^n} h(x) p_{\Sigma_t}(x) dx,$$

is an increasing function of t , $0 \leq t \leq 1$.

Proof: Assuming, as before, that $p(\lambda)$ is a C^2 function with compact support, then (6) and (8) give

$$\frac{\partial}{\partial t} \int_{\mathbb{R}^n} h(x) p_{\Sigma_t}(x) dx = \int_{\mathbb{R}^n} Ah(x) G_{\Sigma_t}(x) dx.$$

By (9) this is non-negative.

References

- Das Gupta, S., Eaton, M.L., Olkin, I., Perlman, M., Savage, L.J. and Sobel, M.
(1972). Inequalities on the probability content of convex regions for elliptically contoured distributions. Proc. Sixth Berkeley Symp. Math. Statist. Prob. 2, 241-264. Univ. of California Press.
- Jogdeo, K. and Proschan, F., (1981). Negative association of random variables, with applications. Florida State Univ. statistics preprint.
- Pitt, L.D., (1981). Positively correlated normal variables are associated. Ann. Prob., to appear.

Plackett, R.L., (1954). A reduction formula for normal multivariate integrals.

Biometrika. 41, 351-360.

Slepian, D., (1962). The one-sided barrier problem for Gaussian noise. Bell

System Tech. J. 41, 463-501.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER AFOSR-TR- 82-0899	2. GOVT ACCESSION NO. AD-A120	3. RECIPIENT'S CATALOG NUMBER 395	
4. TITLE (and Subtitle) ASSOCIATION OF NORMAL RANDOM VARIABLES AND SLEPIAN'S INEQUALITY		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL	
		6. PERFORMING ORG. REPORT NUMBER FSU #M620	
7. AUTHOR(s) Kumar Jogdeo*, Michael D. Perlman**, and *** Loren D. Pitt		8. CONTRACT OR GRANT NUMBER(s) F49620-82-K-0007	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics & Statistical Consulting Center, Florida State University, Tallahassee FL 32306		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PE61102F; 2304/A5	
11. CONTROLLING OFFICE NAME AND ADDRESS Directorate of Mathematical & Information Sciences Air Force Office of Scientific Research Bolling AFB DC 20332		12. REPORT DATE 1982	
		13. NUMBER OF PAGES 8	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Normal variables; association; Slepian's inequality.			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A simple extension of Slepian's inequality is given which implies the original inequality and the result that positively correlated normal variables are associated.			

82 10 18 046

DD FORM 1 JAN 73 1473

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)